

3D Simulations of Galactic Winds in Dwarf Galaxies

Andrea Marcolini and Fabrizio Brighenti

Department of Astronomy, University of Bologna

Annibale D’Ercole

Bologna Astronomical Observatory

Galactic winds play a key role in the formation and evolution of galaxies and the intergalactic medium (IGM). The fate of the heavy elements ejected by the SNII produced in a starbursting dwarf galaxy is a particular crucial issue: are the metals blown out in the IGM and lost forever from the galaxy? Or are they confined by relatively large pressure gas existing at large radii (both ISM or ICM)? In the latter case a significant fraction of the metals might fall back into the galaxy, enriching the ISM. In the simulation presented here we include a new ingredient affecting the evolution of galactic winds: the ram pressure applied by the IGM on the ISM as the galaxy roams through it. We consider a typical dwarf galaxy moving inside a typical poor galaxy group. The model galaxy is built along the lines of D’Ercole & Brighenti (1999, MNRAS, 309, 941), with a stellar disk with mass $M_* \sim 6 \times 10^8 M_\odot$ and a spherical dark halo $M_h \sim 7 \times 10^9 M_\odot$. The ISM, in rotational equilibrium in the resulting potential, has mass $M_g \sim 6 \times 10^8 M_\odot$ in the “galactic region”, defined as $|z| < 1$ kpc, $R < 2$ kpc, in cylindrical coordinates. The galaxy moves with a velocity $v = 200 \text{ km s}^{-1}$ through an ICM with $n_e = 10^{-4} \text{ cm}^{-3}$ and $T = 10^6 \text{ K}$. After 1 Gyr, a central starburst generates a power $L = 3.8 \times 10^{40} \text{ erg s}^{-1}$, for 30 Myr. We follow the evolution of the resulting galactic wind for 500 Myr.

Figure 1 shows the density maps in the equatorial and meridional plane at four different times. The left panel refers to the time just before the starburst onset. The ram pressure limits the radial extent of the




Figure 1. Density maps on the equatorial (bottom row) and meridional (top row) plane, at four different times. Axis labels are in kpc.



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.




Figure 2. Variation of the gas mass (left) and the metal-rich ejecta mass (right) vs. time. Solid line are for the galaxy subject to ram pressure stripping, dashed lines represent the galaxy at rest in the ICM. The total amount of ejecta produced by the burst after 30 Myr is $9 \times 10^5 M_{\odot}$.

ISM to $r \approx 5 - 10$ kpc and generates the classic gas tail. The mass in the galactic region, however, has not been eroded by the stripping.

The other panels of Figure 1 shows the density at the end of the energy injection phase (30 Myr), at $t = 50$ Myr, and at a late time $t = 200$ Myr. The superbubble breaks out injecting the metal-rich gas in the ICM. We accurately follow the evolution of the SN ejecta as a separate fluid. In Figure 2 are plotted the time (0-500 Myr) variations of the ISM mass and the SN ejecta fraction in the galactic region. We find that the starburst is inefficient in removing the ISM. The gas mass slightly decreases for $0 < t < 100$ Myr, then increases again and actually more gas is present at the final time than at the time of the starburst onset. Shortly after the SN activity ceased, a radial flow fills the central hole, the initial ISM distribution is approximately recovered, and the galaxy might be ready for a second starburst.

The SN ejecta behaves differently. At any time only a minor fraction of the metal-rich material is present in the galactic region. Some ejecta flows back in with the ISM flow when the superbubble collapse, but at the end of the simulation only $\sim 3\%$ of the total metals are found inside the galaxy. In the model presented here the ram pressure has little effect on the evolution of the ISM and metal-rich material in the galactic central region (see Figure 2). While the ejecta is carried downstream at large distance by the moving ICM, for a galaxy at rest most of the ejecta is dispersed in the ICM at large heights (~ 30 kpc) over the galaxy. For larger ram pressures, or smaller galaxies, the ISM is significantly modified (or completely removed) by the interaction with the ICM. When this happens, galactic winds become more efficient in blowing out ISM and metals, and a relatively weak starburst may disrupt the ISM of the parent galaxy. A large grid of ram pressure+wind models will be described in detail in Marcolini et al. (in preparation).